
Principles and Practice of Clinical Electrophysiology of Vision

Editors

JOHN R. HECKENLIVELY, M.D.
Professor of Ophthalmology
Jules Stein Eye Institute
Los Angeles, California

GEOFFREY B. ARDEN, M.D., PH.D.
Professor of Ophthalmology and
Neurophysiology
Institute of Ophthalmology
Moorfields Eye Hospital
London, England

Associate Editors

EMIKO ADACHI-USAMI, M.D.
Professor of Ophthalmology
Chiba University School of Medicine
Chiba, Japan

G.F.A. HARDING, PH.D.
Professor of Neurosciences
Department of Vision Sciences
Aston University
Birmingham, England

SVEN ERIK NILSSON, M.D., PH.D.
Professor of Ophthalmology
University of Linköping
Linköping, Sweden

RICHARD G. WELEBER, M.D.
Professor of Ophthalmology
University of Oregon Health Science Center
Portland, Oregon

 **Mosby
Year Book**

St. Louis Baltimore Boston Chicago London Philadelphia Sydney Toronto



Dedicated to Publishing Excellence

Sponsoring Editor: David K. Marshall
Assistant Director, Manuscript Services: Frances M. Perveiler
Production Project Coordinator: Karen E. Halm
Proofroom Manager: Barbara Kelly

Copyright © 1991 by Mosby-Year Book, Inc.
A Year Book Medical Publishers imprint of Mosby-Year Book, Inc.

Mosby-Year Book, Inc.
11830 Westline Industrial Drive
St. Louis, MO 63146

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior written permission from the publisher. Printed in the United States of America.

Permission to photocopy or reproduce solely for internal or personal use is permitted for libraries or other users registered with the Copyright Clearance Center, provided that the base fee of \$4.00 per chapter plus \$.10 per page is paid directly to the Copyright Clearance Center, 21 Congress Street, Salem, MA 01970. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collected works, or for resale.

1 2 3 4 5 6 7 8 9 0 CL CL MV 95 94 93 92 91

Library of Congress Cataloging-in-Publication Data

Principles and practice of visual electrophysiology / [edited by] John R. Heckenlively, Geoffrey B. Arden.

p. cm.

Includes bibliographical references.

Includes index.

ISBN 0-8151-4290-0

1. Electroretinography. 2. Electrooculography. 3. Visual evoked response. I. Heckenlively, John R. II. Arden, Geoffrey B. (Geoffrey Bernard)

[DNLM: 1. Electrooculography. 2. Electrophysiology. 3. Electroretinography. 4. Evoked Potentials, Visual. 5. Vision

Disorders—physiopathology. WW 270 P957]

RE79.E4P75 1991

617.7 1547—dc20

DNLM/DLC

for Library of Congress

91-13378

CIP

Retinal Extracellular Potential Responses Not Evoked by Light

Chester Karwoski

POTENTIALS EVOKED BY OPTIC NERVE STIMULATION

The contribution of ganglion cells to the electroretinogram (ERG) has been explored by antidromically activating them via electrical stimulation of the optic nerve. The first to do this were Ogden and Brown,^{22, 23} who found in primate retina a field potential that had maximum positive amplitude in the inner plexiform layer and a smaller negativity in the ganglion cell layer. This response, termed the *P-wave*, was also observed in primates by Gouras,⁷ but he found a reversal point in the inner plexiform layer and the largest positivity in the inner nuclear layer. A similar response profile was found by Miller et al.¹⁷ in the mudpuppy.

The exact origin of the P-wave is controversial. It was initially thought to arise from activation of inner retinal neurons via the efferent system,²³ but this remains a possibility only in nonmammalian species where retinal efferent innervation has been clearly shown. More recently, arguments have been marshaled for the P-wave arising directly from ganglion cell spikes,⁷ from Müller cell spatial buffering of K⁺ released by ganglion cells,¹⁷ and from activity in retinal neurons that are postsynaptic to ganglion cells.^{22, 28} Other mechanisms may also exist by which activity in ganglion cell axons may induce responses elsewhere in the retina, eg, "blue arcs."² It is possible that multiple mechanisms may contribute to the P-wave, and the size of the contribution may depend on the species and experimental methodology. At any rate, it is not certain whether the P-wave arises *directly* from ganglion cell activity.

There is also disagreement whether the P-wave can be recorded transretinally. Studies in primates failed to find such a contribution, but a substantial negativity in the vitreous was seen in the mudpuppy.¹⁷ As with the proximal negative response (PNR) and M-wave, a contribution from the generators of the P-wave to the transretinal ERG has not been ruled out completely, and a possible role in the pattern ERG should be considered.

POTENTIALS EVOKED BY TRANSRETINAL ELECTRICAL CURRENT

Simultaneous electrical stimulation of a single retinal neuron while recording from another is a powerful technique that has been used by several laboratories to elucidate connections between neurons of the proximal retina.²¹ In addition, single photoreceptors have been stimulated electrically in order to bypass the visual transduction process.¹ Electric current passed extracellularly across the whole retina (transretinal current) will also generate retinal activity that does not arise via phototransduction. The first to demonstrate this was Motokawa et al.,²⁰ who passed current across the isolated retina of carp and recorded an electrically evoked ERG (the EERG). The EERG has since been shown to be a general feature of the vertebrate retina.^{13, 23}

The EERG has several components, and its analysis will likely prove as complex as the standard light-evoked ERG. In the primate, Ogden and Brown²³ presented evidence that three EERG components arise via current-evoked antidromic activation of

ganglion cells, while two others (early negative wave [EN] and late negative wave [LN]), arise in the inner nuclear layer. The EERG LN and ERG b-wave have nearly identical intraretinal depth profiles, and thus these field potentials may arise through the same underlying processes.^{14, 20, 23} However, a comparison of changes in extracellular $[K^+]_o$ during transretinal current and during the light-evoked b-wave does not clearly support this idea.^{11, 12}

Another issue concerns which cells are activated by transretinal current. There is widespread agreement that photoreceptor synaptic terminals are activated^{13, 31, 32} and that activity initiated by this process will continue through the rest of the retinal circuitry. Transretinal current will directly activate bipolar-amacrine synapses in the absence of photoreceptor synaptic transmission³¹ and even in the absence of photoreceptors.²⁶ In addition, ganglion cell axons may be activated directly.²³ Direct electrical activation of other retinal cells and synapses should also be considered.

Because the proximal retina can be directly activated by electrical current, the EERG has potential value as a tool for assessing the integrity of proximal retinal neurons during disorders in which photoreceptors have degenerated or are inactive.²⁵ It may also be useful during disorders in which light cannot reach the photoreceptors (e.g., corneal burns, cataracts, intraocular hemorrhage).

SPREADING DEPRESSION

The phenomenon of spreading depression (SD), first described by Leão¹⁵ in cerebral cortex, consists of a slowly propagating and transient depression of neural activity, and it is associated with a host of physiological events, including a large, local field potential. Gouras⁸ described SD in amphibian retinas and it has since been observed in many species, including mammals.^{16, 27} The field potential associated with retinal SD has maximum amplitude in the proximal retina, possibly at the inner limiting membrane.¹⁹ Within the proximal retina, this potential is negative going, but the transretinally recorded SD potential can be predominately negative or positive, depending on the species.⁵ SD is also associated with a large increase in $[K^+]_o$ in the inner plexiform layer, and several lines of evidence suggest that the origin of the SD potential is currents associated with the spatial buffering of this increased $[K^+]_o$ by Müller cells.^{18, 19, 24}

The frequency of retinal SD can be increased by events such as injury to a retinal region,⁸ mechanical

stimulation of the retina,⁴ anoxia,⁸ application of high concentrations of K ions¹⁰ or low concentrations of Cl ions,^{18, 24} application of the excitatory amino acids glutamate or aspartate,^{18, 29} light offset,²⁷ and antidromic stimulation of the optic nerve¹⁸; seemingly spontaneous SDs have been observed in chick (R.H. Steinberg, personal communication) and lizard eyecups (C.J. Karwoski, unpublished observation). Because of the large number of conditions that facilitate or initiate SDs it seems reasonable that SD will sometimes be associated with disorders of the eye or retina in humans. Although this has not been reported, it is a fruitful topic for future research.²⁷ Two difficulties are that (1) the retinal SD may be a transient event, and the clinician must be recording an ERG or looking in the eye (for the color change of the retina associated with SD³³) at the proper time; (2) it is not certain whether a retinal SD could be detected by electroretinography. In vitreal recordings from the eyecup of the lizard *Anolis*, SDs generate transient positive shifts in potential, following which ERG amplitude is depressed for several minutes (unpublished observation). However, the lizard eye is small, and SD could dominate the recording. An SD in the large eye of humans may depress a relatively small retinal area, and both the SD-induced potential shift and the depression in the ERG that is evoked by diffuse light may be undetectable.

In the brain, increasing evidence points to SD generating many of the symptoms of migraine headaches,^{6, 9} and SD occurs in conjunction with stereotaxic surgery.³ Somjen and Aitkin³⁰ argue for SD to be considered a transient and propagating type of cortical depression that is initiated at one point, while another type of depression is diffuse and non-propagating. The latter would possibly include depressions arising during transient cerebral hypoxia (due to asphyxia or to brief failure of cerebral circulation) and during concussions. Because of the number of events possibly related to cortical depressions (both spreading and diffuse) in humans, the clinician should keep alert to the possibility of such depressions in the retina.

Acknowledgment

I thank Dr. R.W. Knighton for helpful discussion.

REFERENCES

1. Baylor DA, Fettiplace R: Transmission from photoreceptors to ganglion cells in turtle retina. *J Physiol* 1977; 271:391-424.

2. Brindley GS: *Physiology of the Retina and Visual Pathway*. London, Edward Arnold Publishers, Ltd, 1970.
3. Bures J, Buresova O, Krivanek J: The meaning and significance of Leao's spreading depression. *An Acad Bras Cienc* 1984; 56:385–400.
4. DeAzeredo FAM, Martins-Ferreira H: Changes in fluid compartments and ionic composition in the isolated chick retina during SD. *Neurochem Res* 1979; 4:99–107.
5. Fujimoto M, Tomita T: The spreading depression potential and the ERG b-wave in retinas of the frog and chick. *Neurosci Res* 1986; Suppl 4:S197–S208.
6. Gardner-Medwin AR, Mutch WAC: Experiments on spreading depression in relation to migraine and neurosurgery. *An Acad Bras Cienc* 1984; 56:423–430.
7. Gouras P: Antidromic responses of orthodromically identified ganglion cells in monkey retina. *J Physiol* 1969; 240:407–419.
8. Gouras P: Electric activity of toad retina. *Am J Ophthalmol* 1958; 46:59–70.
9. Hansen AJ, Lauritzen M: The role of spreading depression in acute brain disorders. *Ann Acad Bras Cienc* 1984; 56:457–480.
10. Higashida H, Sakakibara M, Mitarai G: Spreading depression in isolated carp retina. *Brain Res* 1977; 120:67–83.
11. Karwoski CJ: Changes in $[K^+]_o$ induced by transretinal currents in frog retina. *Physiol. Bohemoslav* 1988; 37:217–225.
12. Karwoski CJ, Coles JA, Lu H-K, Huang B: Current-evoked transcellular K^+ flux in frog retina. *J Neurophysiol* 1989; 61:939–952.
13. Knighton RW: An electrically evoked slow potential of the frog's retina. I. Properties of response. *J Neurophysiol* 1975; 38:185–197.
14. Knighton RW: An electrically evoked slow potential of the frog's retina. II. Identification with PII component of electroretinogram. *J Neurophysiol* 1975; 38:198–209.
15. Leao AAP: Spreading depression of activity in the cerebral cortex. *J Neurophysiol* 1944; 7:359–390.
16. Miller RF, Dacheux R: Chloride-sensitive receptive field mechanisms in the isolated retina-eyecup of the rabbit. *Brain Res* 1975; 90:329–334.
17. Miller RF, Dacheux R, Proenza LM: Müller cell depolarization evoked by antidromic optic nerve stimulation. *Brain Res* 1977; 121:162–166.
18. Mori S, Miller WH, Tomita T: Microelectrode study of spreading depression (SD) in frog retina—General observations of field potentials associated with SD. *Jpn J Physiol* 1977; 26:203–217.
19. Mori S, Miller WH, Tomita T: Microelectrode study of spreading depression (SD) in frog retina—Müller cell activity and $[K^+]_o$ during SD. *Jpn J Physiol* 1977; 26:219–233.
20. Motokawa K, Yamashita E, Ogawa T: Responses of retinal network to electrical stimulation. *Tohoku J Exp Med* 1959; 71:41–53.
21. Naka K-I: Functional organization of catfish retina. *J Neurophysiol* 1977; 40:26–43.
22. Ogden TE: Intraretinal slow potentials evoked by brain stimulation in the primate. *J Neurophysiol* 1966; 29:898–908.
23. Ogden TE, Brown KT: Intraretinal responses of the cynomolgus monkey to electrical stimulation of the optic nerve and retina. *J Neurophysiol Jpn* 1964; 27:682–705.
24. Olsen JS, Miller RF: Spontaneous slow potentials and spreading depression in amphibian retina. *J Neurophysiol* 1977; 40:752–767.
25. Potts AM, Inoue J: The electrically evoked response (EER) of the visual system. II. Effect of adaptation and retinitis pigmentosa. *Invest Ophthalmol* 1969; 8:605–612.
26. Potts AM, Inoue J: The electrically evoked response of the visual system (EER). III. Further contribution to the origin of the EER. *Invest Ophthalmol* 1970; 9:814–819.
27. Ripps H, Mehaffey L, Siegel IM: "Rapid regeneration" in the cat retina. *J Gen Physiol* 1981; 77:335–346.
28. Sakai HM, Naka K-I, Dowling JE: Ganglion cell dendrites are presynaptic in catfish retina. *Nature* 1986; 319:495–497.
29. Shimazaki H, Karwoski CJ, Proenza LM: Aspartate-induced dissociation of proximal from distal retinal activity in the mudpuppy. *Vision Res* 1984; 24:587–595.
30. Somjen GG, Aitkin PG: The ionic and metabolic responses associated with neuronal depression of Leao's type in cerebral cortex and in hippocampal formation. *An Acad Bras Cienc* 1984; 56:495–504.
31. Toyoda J-I, Fujimoto M: Application of transretinal current stimulation for the study of bipolar-amacrine transmission. *J Gen Physiol* 1984; 84:915–925.
32. Trifinov YA: Study of synaptic transmission between the photoreceptor and the horizontal cell using electrical stimulation of the retina. *Biofizika* 1968; 13:809–817.
33. Van Harrevelde A: Visual concomitants of retinal spreading depression. *An Acad Bras Cienc* 1984; 56:519–524.